



1412 SW 43rd Street · Suite 120

Renton, WA 98057

• 877-425-6337 PHONE

• 877-509-6337 E-Fax

www.readymedspharmacy.com

readymedspharmacy@gmail.com

www.facebook.com/readymedspharmacy

If you are interested in Ready Meds Pharmacy administering flu/pneumonia vaccines at your facility, please return this letter with the attached informed consent completed for each client via fax within 2 weeks. Flu/Pneumonia shots are available through Original Medicare at no charge. We can bill private insurances or Medicare Advantage Plans but there may be an applicable co-pay. Please check with your client's pharmacy insurance carrier if you have any questions on his or her copay.

If your client does not have insurance or his/her insurance does not cover flu vaccination, our prices are as followed:

- Flu shots are \$45.00
- High Dose flu shots are \$75.00
- Pneumonia shots are \$135.00
- Prevnar shots are \$230.00

We have a limited amount of High Dose flu shots available this year, please inquire about availability. High Dose flu shots are first come, first serve.

Our Suppliers do not carry "Preservative Free" shots.

We cannot bill Original Medicare or Medicare Advantages plans for Hospice patients. Claims for hospice patients previous years were all denied. If you have a hospice patient, the only form of payment we can accept is cash.

CDC currently recommends people 65 and over get both Pneumovax 23 and Prevnar 13 vaccines. Some patients under 65 with certain medical conditions may qualify. Please inquire with the pharmacy. Please only check one pneumonia box as both vaccines cannot be given at the same time. If your clients are 65 and over and have ever received either of these pneumonia vaccines, they no longer need the pneumonia shots per CDC recommendations.

Please have payment ready on the day of your flu shot visitation. You may also prepay by calling our pharmacy up to 1 day prior to your visitation. We are not able to include the cost of vaccination on your client's AR account with the pharmacy.

Because of the high volume of requests we get from our adult family homes, we will only be able to visit each home once this flu season. We will try our best to schedule your visitation on a day where all your clients are present. If for any reason one or more of your clients are not available on your visitation date, they will need to find another way to get their vaccine. We are sorry for any inconvenience this may cause you.

Name of Adult Family Home or Facility

Address

Phone Number

Contact Name



1412 SW 43rd Street · Suite 120

Renton, WA 98057

· 877-425-6337 PHONE

· 877-509-6337 E-Fax

www.readymedspharmacy.com

readymedspharmacy@gmail.com

www.facebook.com/readymedspharmacy

Date: _____

Flu Shot High Dose Flu Prevnar 13 Pneumovax 23

Medicare # Other Insurance (provide copy of card) Private Pay (have cash/check ready)

Name: _____ DOB: _____ Sex: M F

Address: _____ Phone # _____

Allergies: _____ Weight (if less than 110lbs) _____

Medical Conditions: _____

Your Primary Physician (and Phone# if known): _____

The following questions will help us determine which vaccines you may be given today. If you answer "YES" to any questions, it does not necessarily mean you should not be vaccinated. It just means additional questions must be asked. If a question is not clear, please ask your healthcare provider to explain it.

	YES	NO	DON'T KNOW
1. Are you sick today?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Do you have allergies to medications, food or any vaccine?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. Have you ever had a serious reaction after receiving a vaccination?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. Do you have a long-term health problem with heart disease, lung disease, asthma, kidney disease, metabolic disease (e.g. diabetes), anemia, or other blood disorder?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. Do you have cancer, leukemia, AIDS, or any other immune system problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6. Do you take cortisone, prednisone, other steroids, or anticancer drugs, or have you had radiation treatments?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7. Have you had a seizure, brain, or other nervous system problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8. During the past year, have you received a transfusion of blood or blood products, or been given immune (gamma) globulin or an antiviral drug?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9. For women: Are you pregnant or is there a chance you could become pregnant during the next month?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
10. Have you received any vaccinations in the past 4 weeks?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Please read the following statements and sign below on the signature line.

I have read or have had explained the information provided about the vaccine I am about to receive. I have received and read a vaccine information statement. I have had a chance to ask questions that were answered to my satisfaction. I believe I understand the benefits and risks of vaccination and ask that the vaccine be given to me or to the person named above for whom I am authorized to make this request. Medicare, I do hereby authorize Ready Meds Pharmacy to release information and request payment. I certify that the information given by me in applying for payment under Medicare is correct. I authorize release of all records to act on this request. I request that payment of authorized benefits be made on my behalf.

X _____ Date: _____

Signature of person to receive vaccine or person authorized to make request (parent or guardian)

For office use only

Vaccine	Vaccine	Vaccine
Lot: #	Lot: #	Lot: #
Mfr.	Mfr.	Mfr.
Date on VIS	Date on VIS	Date on VIS
Site	Site	Site
Date given	Date given	Date given

X _____ Date: _____

Signature of Administrator



1412 SW 43rd Street · Suite 120

Renton, WA 98057

• 877-425-6337 PHONE

• 877-509-6337 E-Fax

www.readymedspharmacy.com

readymedspharmacy@gmail.com

www.facebook.com/readymedspharmacy

Risk Group	Underlying Medical Condition	PCV13		PPSV23	
		Recommended	Recommended	Recommended	Revaccination at 5 years after first dose
Immunocompetent persons	Chronic Heart disease			✓	
	Chronic Lung disease			✓	
	Diabetes mellitus			✓	
	CSF leaks	✓		✓	
	Cochlear implants	✓		✓	
	Alcoholism			✓	
	Chronic liver disease			✓	
	Cigarette smoking			✓	
Persons with functional or anatomic asplenia	Sickle cell disease/other hemoglobinopathies	✓		✓	✓
	Congenital or acquired asplenia	✓		✓	✓
Immunocompromised Persons	Congenital or acquired immunodeficiencies	✓		✓	✓
	HIV infection	✓		✓	✓
	Chronic renal failure	✓		✓	✓
	Nephrotic syndrome	✓		✓	✓
	Leukemia	✓		✓	✓
	Lymphoma	✓		✓	✓
	Hodgkin disease	✓		✓	✓
	Generalized malignancy	✓		✓	✓
	Solid organ transplant	✓		✓	✓
	Multiple myeloma	✓		✓	✓

Influenza (Flu) Vaccine (Inactivated or Recombinant): *What you need to know*

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Influenza (“flu”) is a contagious disease that spreads around the United States every year, usually between October and May.

Flu is caused by influenza viruses, and is spread mainly by coughing, sneezing, and close contact.

Anyone can get flu. Flu strikes suddenly and can last several days. Symptoms vary by age, but can include:

- fever/chills
- sore throat
- muscle aches
- fatigue
- cough
- headache
- runny or stuffy nose

Flu can also lead to pneumonia and blood infections, and cause diarrhea and seizures in children. If you have a medical condition, such as heart or lung disease, flu can make it worse.

Flu is more dangerous for some people. Infants and young children, people 65 years of age and older, pregnant women, and people with certain health conditions or a weakened immune system are at greatest risk.

Each year **thousands of people in the United States die from flu**, and many more are hospitalized.

Flu vaccine can:

- keep you from getting flu,
- make flu less severe if you do get it, and
- keep you from spreading flu to your family and other people.

2 Inactivated and recombinant flu vaccines

A dose of flu vaccine is recommended every flu season. Children 6 months through 8 years of age may need two doses during the same flu season. Everyone else needs only one dose each flu season.

Some inactivated flu vaccines contain a very small amount of a mercury-based preservative called thimerosal. Studies have not shown thimerosal in vaccines to be harmful, but flu vaccines that do not contain thimerosal are available.

There is no live flu virus in flu shots. **They cannot cause the flu.**

There are many flu viruses, and they are always changing. Each year a new flu vaccine is made to protect against three or four viruses that are likely to cause disease in the upcoming flu season. But even when the vaccine doesn’t exactly match these viruses, it may still provide some protection.

Flu vaccine cannot prevent:

- flu that is caused by a virus not covered by the vaccine, or
- illnesses that look like flu but are not.

It takes about 2 weeks for protection to develop after vaccination, and protection lasts through the flu season.

3 Some people should not get this vaccine

Tell the person who is giving you the vaccine:

- **If you have any severe, life-threatening allergies.**

If you ever had a life-threatening allergic reaction after a dose of flu vaccine, or have a severe allergy to any part of this vaccine, you may be advised not to get vaccinated. Most, but not all, types of flu vaccine contain a small amount of egg protein.

- **If you ever had Guillain-Barré Syndrome (also called GBS).**

Some people with a history of GBS should not get this vaccine. This should be discussed with your doctor.

- **If you are not feeling well.**

It is usually okay to get flu vaccine when you have a mild illness, but you might be asked to come back when you feel better.



4 Risks of a vaccine reaction

With any medicine, including vaccines, there is a chance of reactions. These are usually mild and go away on their own, but serious reactions are also possible.

Most people who get a flu shot do not have any problems with it.

Minor problems following a flu shot include:

- soreness, redness, or swelling where the shot was given
- hoarseness
- sore, red or itchy eyes
- cough
- fever
- aches
- headache
- itching
- fatigue

If these problems occur, they usually begin soon after the shot and last 1 or 2 days.

More serious problems following a flu shot can include the following:

- There may be a small increased risk of Guillain-Barré Syndrome (GBS) after inactivated flu vaccine. This risk has been estimated at 1 or 2 additional cases per million people vaccinated. This is much lower than the risk of severe complications from flu, which can be prevented by flu vaccine.
- Young children who get the flu shot along with pneumococcal vaccine (PCV13) and/or DTaP vaccine at the same time might be slightly more likely to have a seizure caused by fever. Ask your doctor for more information. Tell your doctor if a child who is getting flu vaccine has ever had a seizure.

Problems that could happen after any injected vaccine:

- People sometimes faint after a medical procedure, including vaccination. Sitting or lying down for about 15 minutes can help prevent fainting, and injuries caused by a fall. Tell your doctor if you feel dizzy, or have vision changes or ringing in the ears.
- Some people get severe pain in the shoulder and have difficulty moving the arm where a shot was given. This happens very rarely.
- Any medication can cause a severe allergic reaction. Such reactions from a vaccine are very rare, estimated at about 1 in a million doses, and would happen within a few minutes to a few hours after the vaccination.

As with any medicine, there is a very remote chance of a vaccine causing a serious injury or death.

The safety of vaccines is always being monitored. For more information, visit: www.cdc.gov/vaccinesafety/

5 What if there is a serious reaction?

What should I look for?

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or unusual behavior.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would start a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 and get the person to the nearest hospital. Otherwise, call your doctor.
- Reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor should file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS does not give medical advice.

6 The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling **1-800-338-2382** or visiting the VICP website at www.hrsa.gov/vaccinecompensation. There is a time limit to file a claim for compensation.

7 How can I learn more?

- Ask your healthcare provider. He or she can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/flu

Vaccine Information Statement
Inactivated Influenza Vaccine

08/07/2015

42 U.S.C. § 300aa-26

Office Use Only



Pneumococcal Conjugate Vaccine (PCV13)

What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Vaccination can protect both children and adults from **pneumococcal disease**.

Pneumococcal disease is caused by bacteria that can spread from person to person through close contact. It can cause ear infections, and it can also lead to more serious infections of the:

- Lungs (pneumonia),
- Blood (bacteremia), and
- Covering of the brain and spinal cord (meningitis).

Pneumococcal pneumonia is most common among adults. Pneumococcal meningitis can cause deafness and brain damage, and it kills about 1 child in 10 who get it.

Anyone can get pneumococcal disease, but children under 2 years of age and adults 65 years and older, people with certain medical conditions, and cigarette smokers are at the highest risk.

Before there was a vaccine, the United States saw:

- more than 700 cases of meningitis,
- about 13,000 blood infections,
- about 5 million ear infections, and
- about 200 deaths

in children under 5 each year from pneumococcal disease. Since vaccine became available, severe pneumococcal disease in these children has fallen by 88%.

About 18,000 older adults die of pneumococcal disease each year in the United States.

Treatment of pneumococcal infections with penicillin and other drugs is not as effective as it used to be, because some strains of the disease have become resistant to these drugs. This makes prevention of the disease, through vaccination, even more important.

2 PCV13 vaccine

Pneumococcal conjugate vaccine (called PCV13) protects against 13 types of pneumococcal bacteria.

PCV13 is routinely given to children at 2, 4, 6, and 12–15 months of age. It is also recommended for children and adults 2 to 64 years of age with certain health conditions, and for all adults 65 years of age and older. Your doctor can give you details.

3 Some people should not get this vaccine

Anyone who has ever had a life-threatening allergic reaction to a dose of this vaccine, to an earlier pneumococcal vaccine called PCV7, or to any vaccine containing diphtheria toxoid (for example, DTaP), should not get PCV13.

Anyone with a severe allergy to any component of PCV13 should not get the vaccine. *Tell your doctor if the person being vaccinated has any severe allergies.*

If the person scheduled for vaccination is not feeling well, your healthcare provider might decide to reschedule the shot on another day.

4 Risks of a vaccine reaction

With any medicine, including vaccines, there is a chance of reactions. These are usually mild and go away on their own, but serious reactions are also possible.

Problems reported following PCV13 varied by age and dose in the series. The most common problems reported among children were:

- About half became drowsy after the shot, had a temporary loss of appetite, or had redness or tenderness where the shot was given.
- About 1 out of 3 had swelling where the shot was given.
- About 1 out of 3 had a mild fever, and about 1 in 20 had a fever over 102.2°F.
- Up to about 8 out of 10 became fussy or irritable.

Adults have reported pain, redness, and swelling where the shot was given; also mild fever, fatigue, headache, chills, or muscle pain.

Young children who get PCV13 along with inactivated flu vaccine at the same time may be at increased risk for seizures caused by fever. Ask your doctor for more information.



Problems that could happen after any vaccine:

- People sometimes faint after a medical procedure, including vaccination. Sitting or lying down for about 15 minutes can help prevent fainting, and injuries caused by a fall. Tell your doctor if you feel dizzy, or have vision changes or ringing in the ears.
- Some older children and adults get severe pain in the shoulder and have difficulty moving the arm where a shot was given. This happens very rarely.
- Any medication can cause a severe allergic reaction. Such reactions from a vaccine are very rare, estimated at about 1 in a million doses, and would happen within a few minutes to a few hours after the vaccination.

As with any medicine, there is a very small chance of a vaccine causing a serious injury or death.

The safety of vaccines is always being monitored. For more information, visit: www.cdc.gov/vaccinesafety/

5

What if there is a serious reaction?

What should I look for?

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or unusual behavior.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness—usually within a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get the person to the nearest hospital. Otherwise, call your doctor.

Reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor should file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS does not give medical advice.

6

The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling **1-800-338-2382** or visiting the VICP website at www.hrsa.gov/vaccinecompensation. There is a time limit to file a claim for compensation.

7

How can I learn more?

- Ask your healthcare provider. He or she can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines

Vaccine Information Statement
PCV13 Vaccine

11/05/2015

42 U.S.C. § 300aa-26

Office Use Only



Pneumococcal Polysaccharide Vaccine

What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Vaccination can protect older adults (and some children and younger adults) from **pneumococcal disease**.

Pneumococcal disease is caused by bacteria that can spread from person to person through close contact. It can cause ear infections, and it can also lead to more serious infections of the:

- Lungs (pneumonia),
- Blood (bacteremia), and
- Covering of the brain and spinal cord (meningitis). Meningitis can cause deafness and brain damage, and it can be fatal.

Anyone can get pneumococcal disease, but children under 2 years of age, people with certain medical conditions, adults over 65 years of age, and cigarette smokers are at the highest risk.

About 18,000 older adults die each year from pneumococcal disease in the United States.

Treatment of pneumococcal infections with penicillin and other drugs used to be more effective. But some strains of the disease have become resistant to these drugs. This makes prevention of the disease, through vaccination, even more important.

2 Pneumococcal polysaccharide vaccine (PPSV23)

Pneumococcal polysaccharide vaccine (PPSV23) protects against 23 types of pneumococcal bacteria. It will not prevent all pneumococcal disease.

PPSV23 is recommended for:

- All adults 65 years of age and older,
- Anyone 2 through 64 years of age with certain long-term health problems,
- Anyone 2 through 64 years of age with a weakened immune system,
- Adults 19 through 64 years of age who smoke cigarettes or have asthma.

Most people need only one dose of PPSV. A second dose is recommended for certain high-risk groups. People 65 and older should get a dose even if they have gotten one or more doses of the vaccine before they turned 65.

Your healthcare provider can give you more information about these recommendations.

Most healthy adults develop protection within 2 to 3 weeks of getting the shot.

3 Some people should not get this vaccine

- Anyone who has had a life-threatening allergic reaction to PPSV should not get another dose.
- Anyone who has a severe allergy to any component of PPSV should not receive it. Tell your provider if you have any severe allergies.
- Anyone who is moderately or severely ill when the shot is scheduled may be asked to wait until they recover before getting the vaccine. Someone with a mild illness can usually be vaccinated.
- Children less than 2 years of age should not receive this vaccine.
- There is no evidence that PPSV is harmful to either a pregnant woman or to her fetus. However, as a precaution, women who need the vaccine should be vaccinated before becoming pregnant, if possible.



4 Risks of a vaccine reaction

With any medicine, including vaccines, there is a chance of side effects. These are usually mild and go away on their own, but serious reactions are also possible.

About half of people who get PPSV have mild side effects, such as redness or pain where the shot is given, which go away within about two days.

Less than 1 out of 100 people develop a fever, muscle aches, or more severe local reactions.

Problems that could happen after any vaccine:

- People sometimes faint after a medical procedure, including vaccination. Sitting or lying down for about 15 minutes can help prevent fainting, and injuries caused by a fall. Tell your doctor if you feel dizzy, or have vision changes or ringing in the ears.
- Some people get severe pain in the shoulder and have difficulty moving the arm where a shot was given. This happens very rarely.
- Any medication can cause a severe allergic reaction. Such reactions from a vaccine are very rare, estimated at about 1 in a million doses, and would happen within a few minutes to a few hours after the vaccination.

As with any medicine, there is a very remote chance of a vaccine causing a serious injury or death.

The safety of vaccines is always being monitored. For more information, visit: www.cdc.gov/vaccinesafety/

5 What if there is a serious reaction?

What should I look for?

Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or unusual behavior.

Signs of a **severe allergic reaction** can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would usually start a few minutes to a few hours after the vaccination.

What should I do?

If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get to the nearest hospital. Otherwise, call your doctor.

Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor might file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS does not give medical advice.

6 How can I learn more?

- Ask your doctor. He or she can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines

Vaccine Information Statement
PPSV Vaccine

4/24/2015

Office Use Only

